

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 October 2003 (23.10.2003)

PCT

(10) International Publication Number
WO 03/086347 A1

(51) International Patent Classification⁷: **A61K 9/10, 9/72**

(21) International Application Number: **PCT/EP03/03916**

(22) International Filing Date: **15 April 2003 (15.04.2003)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
MI2002A000808 17 April 2002 (17.04.2002) IT

(71) Applicant (*for all designated States except US*): **CHIESI FARMACEUTICI S.P.A.** [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **SAMBUCO, Barbara** [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT).
PIRROTTA, Daniele [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT).

(74) Agents: **MINOJA, Fabrizio** et al.; Bianchetti Bracco Minoja S.r.l., Via Rossini, 8, I-20122 Milano (IT).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **PROCESS FOR PREPARATION OF A STERILE SUSPENSION OF CORTICOSTEROID PARTICLES FOR THE ADMINISTRATION BY INHALATION**

(57) Abstract: A process for the preparation of aqueous suspensions of sterile micronized drug particles, in particular corticosteroid, to be administered by inhalation, which produces homogenous dispersions of particles characterised by optimal size and size distribution is disclosed. The process is carried out by using a turboemulsifier equipped with a high-power turbine and connected to a loading hopper.

WO 03/086347 A1

PROCESS FOR PREPARATION OF A STERILE SUSPENSION OF CORTICOSTEROID PARTICLES FOR ADMINISTRATION BY INHALATION

The present invention relates to a process for the preparation of aqueous suspensions of drug particles, to be administered by inhalation, which produces homogenous dispersions of particles characterised by optimal size and size distribution.

PRIOR ART

The method of delivering drugs by inhalation has been used for several years, and is the mainstay of the treatment of diseases that limit the respiratory flow, such as asthma and chronic bronchitis.

The advantages of inhalation over the systemic route include the fact that the drug is released directly at the site of action, thus preventing systemic side effects and resulting in a more rapid clinical response and a higher therapeutic index.

Among the various types of drug which are administered by inhalation for the treatment of the respiratory diseases, corticosteroids, such as beclomethasone dipropionate (BDP), mometasone furoate, flunisolide, budesonide, fluticasone propionate and others are of great importance. They are generally administered in micronised form in suspension, in an aqueous phase that usually also contains surfactants and/or cosolvents, or in a propellant. The drug is inhaled in aerosol form, ie. in the form of a dispersion of solid particles in a gaseous medium. The efficacy of this form of administration depends on the deposit of a sufficient quantity of particles at the site of action.

In order to ensure an effective penetration into the low respiratory tract of the patient, i.e. bronchioli and alveoli, one of the most important parameters

is particle size, which must be equal or lower than 5-6 μm . This size is quantified by measuring a characteristic sphere-equivalent diameter, known as the median aerodynamic diameter (MAD), which expresses the ability of particles to be transported in suspension in an air flow.

5 Particles with a larger MAD are ineffective because they are deposited in the oropharyngeal cavity, and are therefore unable to reach the terminal branches of the respiratory tree; they can also give rise to local side effects, or may be absorbed through the buccal mucosa and give rise to systemic side effects.

10 Another important characteristic to ensure correct administration, and therefore therapeutic efficacy, is homogenous dispersion of the particles in suspension, without the formation of aggregates which prevent correct aerosolisation. The formation of more or less compact aggregates can also give rise to problems of distribution and therefore of uniformity of dose
15 during the filling of the containers. From the technological standpoint it is also very important for the particles to fall within the narrowest and most homogenous possible size distribution range, and to be as finest as possible compatibly with the upper limit (5-6 μm); this because upon variation of environmental humidity conditions, aqueous phase suspensions may face
20 problems over time in terms of constancy of particle distribution due to the total or partial recrystallisation of the small amount of dissolved solute (Davis S *et al* Int J Pharm 1, 303-314, 1978; Tiano S *et al* Pharm Dev Tech 1, 261-268, 1996; Taylor K *et al* Int J Pharm 153, 93-104, 1997). As this parameter is inversely correlated with the MAD of the particles, such an increase can
25 prejudice the efficacy of nebulisation and therapeutic efficacy, as particles with an MAD exceeding 5-6 μm are unable to reach the preferential site of action.

Therefore the finer the particles, the lower the probability that after

partial recrystallisation they will reach the critical size liable to prejudice the properties of the formulation in terms of technological and therapeutic parameters.

Another important requirement that must be met by pharmaceutical formulations for inhalation is sterility. This requirement is becoming more and more mandatory as confirmed by the FDA final rule "*Sterility Requirement for Aqueous-Based Drug Products for Oral Inhalation*" published in the Federal Register of May 26, 2000 (65 FR 34082) governing the quality and safety of pharmaceutical products for a number of reasons, including the fact that the lungs are a particularly vulnerable organ of the human body, and many patients who use inhaled drugs have general health problems.

The current trend is to produce inhalation formulations devoid of preservatives and bacteriostatics, as it has been reported in the literature that some of the substances commonly used for this purpose can induce allergic reactions or give rise to irritation of the respiratory mucosae (Menendez R *et al J Allergy Clin Immunol* 84, 272-274, 1989; Afferty P *et al Thorax* 43, 446-450, 1988). Various processes can be used to manufacture sterile pharmaceutical formulations for inhalation. For example, the active ingredient can be sterilised by dry heating or irradiation, followed by preparation of the formulation under aseptic conditions, or the formulation can be pre-prepared and sterilised by treatment in an autoclave or by filtration.

Some of the sterilisation methods reported suffer from drawbacks or limitations. For example, heat treatments are unsuitable in the case of aqueous suspensions of thermolabile corticosteroids such as beclomethasone dipropionate (BDP), and sterilising filtration is not feasible for suspensions.

WO 99/25359 relates to a process for sterilising corticosteroids by heating them at lower temperatures than those reported in some Pharmacopoeias (110–130°C vs 140-180°C) but does not contain any teaching

as to how to prepare the relevant pharmaceutical formulations in the form of suspensions.

In the patent application WO 00/25746, the applicant described a process for the preparation of aqueous suspensions for nebulisation based on a
5 micronised active ingredient sterilised with gamma rays.

Said process basically involves a first stage of preparation of an aqueous solution, which constitutes the vehicle and contains suitable excipients, in a turboemulsifier, followed by the addition of a sterile micronised active ingredient which in turn is dispersed at atmospheric
10 pressure in the same turboemulsifier. The dispersion of the active ingredient in the aqueous phase may be subjected to an additional high-pressure homogenising treatment which further reduces the average size of the particles in suspension.

In the text an example (Ex. 2) of preparation of a suspension
15 formulation on a pilot scale (100 litres) starting from micronised BDP sterilised by gamma radiation is reported. In said example, the active ingredient is added to the sterile aqueous base and dispersed, initially using magnetic agitation only, then by using a turbine system for 15-20 minutes.

However, when this process has been applied on an industrial scale, it
20 has been found that long processing times are required for the homogenisation stage. A mixing time of over two hours is required for preparations exceeding 1000 litres. Moreover, the obtained dispersions do not meet the requirement of homogeneity in a satisfactory way.

It has been observed that these drawbacks are largely attributable to the
25 technological characteristics of the sterile micronised active ingredient, which disperses more slowly as well as more difficulty in the aqueous vehicle than the unsterilised compound. In fact, sterile micronised corticosteroid particles must be stored under vacuum to maintain their sterility, and consequently tend

to pack more strongly than non-sterile particles of the same active ingredient, as demonstrated by density measurements. Stronger packing is in turn responsible for difficulties with dispersion.

SUMMARY OF THE INVENTION

5 A process for the preparation on an industrial scale of aqueous suspensions for nebulisation comprising a sterile micronised active ingredient, preferably sterilised by irradiation with gamma rays has now been found, and its the object of this invention. The process according to the invention reduces processing times and gives rise to suspensions with a homogenous,
10 reproducible particle distribution and optimum particle-size distribution, thus producing compositions with a high level of physical stability and therapeutic efficacy. Suspensions obtained with the process according to the invention are used as pharmaceutical formulations for aerosol inhalation after being introduced into suitable containers such as multidose vials for nebulisation,
15 and preferably monodose vials.

In the embodiment of the invention, the process is carried out with the use of a turboemulsifier fitted with a high-power turbine, and is characterised in that the active ingredient in powder form is transferred through the turbine by exploiting the vacuum applied in the turboemulsifier. On the contrary, in
20 the prior art, the active ingredient is added from the top directly into the turboemulsifier.

It has now been found that by operating in accordance with the teaching of this invention, i.e. by loading the sterile active ingredient into the turboemulsifier through the turbine after applying the vacuum rather than
25 loading it from the top at atmospheric pressure, far more efficient dispersal of the active ingredient, and therefore homogenous suspensions with a distribution reproducible from one batch to another, can be obtained in a much shorter time, on an industrial scale. By operating under vacuum it is also

possible to prevent foam formation, and therefore the additional operation to remove it. Moreover, it has been unexpectedly found that finer particles with a narrower, more homogenous particle-size distribution range can be obtained by the process of the invention, with no need for further treatments such as
5 treatment in a high-pressure homogeniser as described in WO 00/25746. As already mentioned, these properties give rise to significant advantages during the step of filling the bulk suspension into suitable containers (multidose or monodose vials) and during storage.

In suspensions obtained with the process according to the invention, the
10 particles sediment more slowly because of their finer size, in accordance with Stokes' law, expressed by the formula:

$$V = \frac{d^2 (\rho - \rho_0) g}{18 \eta}$$

15 wherein V is the sedimentation rate, d is the mean diameter of the particles, η is the viscosity of the medium in poises, ρ is the density of the particles, ρ_0 is the density of the dispersing medium, and g is the gravity acceleration.

During the container-filling step, the re-circulation conditions to which
20 the particles of active ingredient are subjected are sufficient to achieve uniform distribution of the particles in the containers, as they pass through the system of radial nozzles of the turbine, and there is no need to use external elements such as pipes or blades to maintain the particles in suspension. The use of such elements would make it necessary to open the apparatus
25 periodically for cleaning operations, thus prejudicing continuity of manufacture under sterile conditions. In the process of the invention, the particles not only sediment more slowly, but also are less liable to form agglomerates, which means that once introduced into the vials, the

suspensions will be more physically stable, and the storage period can be increased. The formation of agglomerates, especially "cakes", ie. sets of highly compact suspended particles, can prejudice the correct dosage of the drug, or at least make administration less therapeutically effective, as the dose
5 may be transferred incompletely from the vial to the bulb of the nebuliser apparatus.

DETAILED DESCRIPTION OF THE INVENTION

The invention will now be described in detail by reference to Figure 1, which shows a scheme of a plant that can be used for the process according to
10 the invention.

A vacuum turboemulsifier (1), constituted by a steel container (2) and fitted with a high-power turbine, and possibly with an agitation system, can be advantageously used to prepare the suspension. "High-power turbine" means a turbine with a power of between 15 to 55 Kwatts. A turboemulsifier, which
15 can agitate the suspension via the system of radial nozzles of the turbine (3), through which the active ingredient passes, will preferably be used, fitted with a 30 Kwatt turbine.

The system is also equipped with a hopper (4) fitted inside an isolator (5) and connected to the turbine of the turboemulsifier via a rigid pipe or hose
20 (6) for the purpose of loading the powder. The entry of the powder into the pipe can be regulated by a butterfly valve to minimise the quantity of incoming air which can contribute to foam formation. "Isolator" means a transparent container, generally made of plexiglas or polyvinylchloride (PVC), fitted with one or more entrance doors and handling gloves for transfer of the
25 powder.

The first stage of the preparation process involves preparing the aqueous solution constituting the vehicle in a suitable tank, preferably made of stainless steel; the solution, which can be sterilised by heat or filtration, may

contain suitable additives or excipients, preferably selected from wetting agents such as polysorbate 20 or sorbitan monolaurate, isotonic agents such as sodium chloride, and optionally stabilising agents such as disodium edetate and/or buffers. The vehicle is preferably sterilised at 121°C for 20 minutes. If
5 necessary, the solution thus obtained is subjected to clarifying filtration and transferred to a turboemulsifier equipped with a vacuum pump. At the second stage, after applying the vacuum in the turboemulsifier, one or more sterile micronised active ingredients are added to the aqueous medium by introducing them from the loading hopper through the turbine.

10 Alternatively, the aqueous solution constituting the vehicle can be prepared and sterilisation performed in a turboemulsifier fitted with a jacket suitable both for steam heating and water cooling.

Advantageously, the active ingredient will be a corticosteroid such as beclomethasone dipropionate, mometasone furoate, fluticasone propionate,
15 flunisolide, ciclesonide or budesonide, micronised by usual processes and sterilised by radiation or heating. Preferably, the active ingredient will be micronised beclomethasone dipropionate sterilised by treatment with gamma rays under the conditions reported in WO 00/25746. At the third stage, the active ingredient is homogenised, again under vacuum, using the turbine
20 system and operating at a speed of between 750 and 4000 rpm, preferably between 1000 and 3600 rpm, and even more preferably between 1600 and 3000 rpm, for 5-60 minutes, and preferably for 20-40 minutes. In the preferred conditions a turbine system operating at 2900 rpm for 30 minutes is used.

25 The suspension of micronised product obtained at the end of the treatment is distributed into suitable containers, preferably constituted by pre-formed monodose vials for nebulisation, optionally pre-sterilised by beta rays irradiation or made with the "blow, fill and seal" technology.

As a consequence, this invention also relates to pharmaceutical formulations to be used for nebulisation, preferably in unit dose preparations containing the aqueous suspensions obtained by the process according to the invention.

5 In said formulations, the Feret diameter of at least 90% of the suspended particles in the final container will advantageously be less than or equal to 8 μm . Preferably, the diameter of at least 50% of the particles will be less than 3 μm , and that of at least 90% less than 7 μm . Even more preferably the diameter of at least 50% of the particles will be less than 2.5 μm , and that
10 of at least 90% less than 6 μm . "Feret diameter" means the distance between imaginary parallel lines tangential to a randomly oriented particle and perpendicular to the ocular scale (USP 26, 2003, page 2185).

The dimensional characteristics of the particles were also evaluated by using a Malvern apparatus. This type of test exploits the diffraction of a laser
15 beam by the particles to determine the size distribution of the particles in suspension. The parameter considered is the median volumetric diameter in μm of 10%, 50% and 90% of the particles, expressed as $d(v,0.1)$, $d(v,0.5)$ and $d(v,0.9)$ respectively, which is determined by assuming that the particles have a geometrical shape equivalent to a sphere. Advantageously, in the suspension
20 formulation of the invention, the $d(v,0.9)$ after sonication is less than 8 μm and the $d(v,0.5)$ is comprised between 2 and 3.5 μm . More preferably the $d(v,0.9)$ is less than 7 μm , the $d(v,0.5)$ is between 2.5 and 3 μm and the particle size distribution (i.e. the difference between $d(v,0.9)$ and $d(v, 0.1)$) does not span for more than 7 μm , preferably for more than 6 μm .

25 The concentration of active ingredient in the pharmaceutical formulations according to the invention is between 0.01 and 0.1% w/v, preferably 0.04% w/v in the case of BDP and 0.025-0.05% in the case of budesonide.

A further object of this invention is the use of the pharmaceutical formulations in unit dose preparations containing the aqueous suspensions obtained by the process according to the invention to treat lung diseases such as asthma and chronic bronchitis with a single daily administration.

5 As reported in the literature for budesonide (Tunek et al. Drug Metab Dispos 1997, 25, 1311-1317), it has been found that stable esters of the active metabolite of BDP, namely beclomethasone-17-monopropionate (17-BMP), are formed in the microsomes of human lung cells with long-chain fatty acids such as oleic acid.

10 These esters are retained in the cells much longer than the parent steroid and release the unchanged active ingredient in a controlled manner, so that the period of tissue exposure to the drug is increased.

It has also been observed that due to the formation of the monoester in the lung cells, the elimination half-life and the mean residence time of 17-BMP are longer after the administration of aqueous suspensions of BDP by
15 nebulisation than formulations in suspension administered in the form of pressurised aerosols .

This last finding has been attributed to the particle distribution obtained after (re)suspension of the micronised active ingredient in the aqueous
20 vehicle. As it can be appreciated from Table 2 and Figure 2 of Example 3 the particles of the sterile micronised active ingredient of the aqueous suspension according to the invention are indeed much finer than those obtained according to the prior art and have a narrower and more homogeneous particle size distribution too. Said particles can more easily dissolve in the lung fluids
25 and penetrate into the cells in a better way, enabling the active ingredient, due to the formation of the esters into the cells, to persist at the site of action for a longer period, so giving rise to a prolonged activity.

As a consequence of the optimal characteristics in term of particle size

of the suspension formulations achieved by the process according to the invention, as well as of the behaviour observed for their active metabolites, pharmaceutical formulations for nebulisation of BDP and budesonide useful for treating lung diseases with a single daily administration can be obtained.

5 This constitutes a considerable advantage in terms of compliance by patients.

The invention is more particularly illustrated in the examples below.

EXAMPLES

10 EXAMPLE 1: Technological characteristics of micronised beclomethasone dipropionate (BDP) sterilised by irradiation with gamma rays compared with the equivalent unsterilised product.

Sterile micronised BDP was obtained as described in WO 00/25746.

The apparent volumes and densities were measured according to the European Pharmacopoeia, 4th edition, paragraph 2.9.15.

15 100 g of the test substance is introduced into a dry 250 ml cylinder without compacting it. The unsettled apparent volume (V_0) is read off, then 10, 500 and 1250 taps are performed, and volumes (V_{10} , V_{500} and V_{1250}) are read. If the difference between V_{500} and V_{1250} is greater than 2 ml, 1250 further taps are performed (V_{2500}).

20 Table 1 shows: i) the apparent density before settling (d_v), which is the ratio between weight (g) and volume before settling (ml); ii) the apparent density after packing (d_s), which is the ratio between weight (g) and the volume after compacting (ml); iii) packing capacity (C_s), which is the difference between V_{10} and V_{500} (ml).

25 The dimensional characteristics of the particles were evaluated by a Malvern apparatus. The results are set out in Table 1.

Table 1 – Technological characteristics of micronised BDP before and after sterilising radiation

Technological characteristics	Non-irradiated BDP	Irradiated BDP
dv (g/ml)	0.21	0.32
ds (g/ml)	0.27	0.42
Cs (ml)	16	12
Particle size (μm , Malvern)		
d (v,0.1)	0.49	0.48
d (v,0.5)	1.91	1.81
d (v,0.9)	5.98	5.73

- 5 The results demonstrate that after radiation, although BDP does not undergo any variations in particle size, it is packed to a greater extent than the non-irradiated product, as indicated by the “packing capacity” value (Cs).

EXAMPLE 2: Preparation of a sterile suspension of micronised BDP sterilised with gamma rays.

10 **Composition:**

Ingredients	Total amount of the preparation	Amount per pharmaceutical unit
Sterile micronised BDP	0.6 kg	(0.8 mg)
Polysorbate (Tween) 20	1.5 kg	(2.0 mg)
15 Sorbitan monolaurate	0.3 kg	(0.4 mg)
Sodium chloride	13.5 kg	(18.0 mg)
Water for injection q. s. for	1500 l	(2.0 ml)

- 20 The first stage of preparation of the sterile suspension involves preparing the aqueous base in a Unimix turboemulsifier fitted with a 30 Kwatt turbine.

After loading water for injection at 60-70°C into the apparatus, sodium chloride and surfactants are added, and the preparation is mixed with the turbine to obtain a homogenous dispersion of the surfactants.

The preparation is then sterilised in a turboemulsifier fitted with a

jacket suitable for both steam heating and water cooling; the sterilisation treatment is conducted at 121°C for approx. 20 minutes.

After filtering and cooling the preparation to the temperature of 30-35°C, a vacuum is applied in the turboemulsifier and the sterile BDP is transferred to the sterile aqueous vehicle through the turbine using the vacuum applied. The active ingredient is dispersed under vacuum along the whole homogenisation stage using the turbine system at 2900 rpm for 30 minutes.

The turboemulsifier is subsequently connected via sterile piping to the storage tank of the container-filling machine and positioned under laminar-flow hood in a controlled-contamination environment, and the suspension is distributed in monodose vials to the volume of 2.15 ml using the "blow, fill and seal" technology.

EXAMPLE 3: Particle-size analysis of preparations obtained according to example 2.

The dimensional characteristics of the particles were evaluated by using a Malvern apparatus and by microscopy.

The Malvern tests were conducted as reported in Example 1. The median volumetric diameter of the particles was determined before and after sonication.

For the purpose of examination under the microscope, a drop of suspension was placed on a slide and covered with a slide cover. The diameter of the particles, expressed as the Feret diameter, was measured with the aid of a micrometer.

The results, expressed as $d(v,0.1)$, $d(v,0.5)$ and $d(v,0.9)$, i.e. as the diameter in μm of 10%, 50% and 90% of the particles, are set out in Table 2, for the purpose of comparison with those relating to a suspension obtained as described in WO 00/25746.

The data relating to the relative distribution frequency of the particle

diameters, measured microscopically, are shown in Figure 2, for suspensions obtained with the process according to the invention (a) and according to the process described in WO 00/25746 (b) respectively.

Table 2: Particle-size characteristics of two sterile suspensions of BDP prepared according to example 2 (Prep. 1) and according to the process described in WO 00/25746 (Prep. 2) respectively.

Particle-size characteristics	Prep. 1 (μm)	Prep. 2 (μm)
Feret Diameter (microscopy)		
d(0.1)	0.35	2.04
d(0.5)	1.82	5.75
d(0.9)	5.18	13.89
Median volumetric diameter (Malvern) without sonication		
d(v,0.1)	0.78	1.32
d(v,0.5)	2.97	6.54
d(v,0.9)	7.88	15.94
Median volumetric diameter (Malvern) after sonication		
d(v,0.1)	0.77	0.96
d(v,0.5)	2.59	4.51
d(v,0.9)	6.25	11.54

The results shown in Table 2 and Figure 2 confirm that the process according to the invention produces finer particles with a narrower and more homogenous particle-size distribution.

CLAIMS

1. A process for the preparation of aqueous suspensions to be used in pharmaceutical formulations for inhalation by nebulisation, which comprises
5 the following steps:
 - a) adding one or more active ingredients in the form of a sterile micronised powder to an aqueous solution constituting the vehicle in a turboemulsifier equipped with a high-power turbine and connected to a loading hopper by introducing said active ingredients through the turbine
10 after having applied the vacuum;
 - b) homogenising the active ingredient in the suspension, still under vacuum, using the turbine system;
 - c) distributing the suspension thus obtained into suitable containers such as monodose vials.
- 15 2. A process as claimed in claim 1, wherein the aqueous solution constituting the vehicle contains additives or excipients selected from wetting, stabilising, isotonic and/or buffer agents.
3. A process as claimed in claims 1 or 2 wherein the active ingredient is a corticosteroid.
- 20 4. A process as claimed in claims 1-3 wherein the aqueous solution constituting the vehicle has been sterilised by heat or filtration.
5. A process as claimed in claims 1-4, wherein the preparation is performed under sterile conditions by adding a micronised corticosteroid, sterilised by irradiation or heating.
- 25 6. A process as claimed in claims 1-5, wherein the corticosteroid is micronised beclomethasone dipropionate sterilised by gamma radiation.
7. A process as claimed in claims 1 to 6, wherein the homogenisation stage is conducted at a speed of between 750 and 4000 rpm for 5 to 60


minutes.

8. A process as claimed in claim 7, wherein the homogenisation stage is conducted at a speed of between 1600 and 3000 rpm for 20 to 40 minutes.
- 5 9. A process as claimed in claim 7, wherein the homogenisation stage is conducted at a speed of 2900 rpm for 30 minutes.
10. A process as claimed in claims 1 to 9, wherein the Feret diameter of at least 90% of the particles is less than or equal to 8 μm .
11. Pharmaceutical formulations to be administered by nebulisation,
10 containing aqueous suspensions obtained with the process claimed in claims 1 to 10.
12. Pharmaceutical formulations as claimed in claim 11, wherein the active ingredient is a corticosteroid selected from BDP, mometasone furoate, flunisolide, budesonide, fluticasone propionate or ciclesonide.
- 15 13. Pharmaceutical formulations as claimed in claim 12, wherein the concentration of active ingredient is between 0.01 and 0.1% w/v.
14. Pharmaceutical formulations as claimed in any one of claims 9 to 13 in unit dose preparations which are pre-formed or produced with the "blow, fill and seal" technology.
- 20 15. Pharmaceutical formulations in the form of aqueous suspension to be administered by nebulization comprising as active ingredient a micronised sterile corticosteroid wherein the median volumetric diameter of the 90% of the particles is less than 8 μm and that of 50% of the particles is comprised between 2 and 3.5 μm as determined by using a
25 Malvern apparatus.
16. Pharmaceutical formulations according to claim 15 wherein the median volumetric diameter of the 90% of the particles is less than 7 μm and that of the 50% of the particles is comprised between 2.5 and 3 μm .

17. Pharmaceutical formulations according to claims 15 and 16 wherein the micronized sterile corticosteroid is beclometasone dipropionate.
 18. Pharmaceutical formulations as claimed in claims 15 to 17, for the treatment of lung diseases such as asthma and chronic bronchitis through an once-a-day administration.
- 5

1/2


Figure 1



2/2


Figure 2 – Particle distribution frequency according to size ranges, expressed by the Feret diameter determined microscopically: a) obtained by the process according to the invention; b) obtained by the process described in WO 00/25746.

Relative frequency %



Feret Diameter (μm)

Relative frequency %



Feret Diameter (μm)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/Eur J3/03916

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K9/10 A61K9/72

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 25746 A (BRAMBILLA GAETANO ;BERNINI EVA (IT); CHIESI PAOLO (IT); MALVOLTI C) 11 May 2000 (2000-05-11) cited in the application page 2, line 13 -page 4, line 16 examples 1,2,6-8 -----	11-18

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search 8 August 2003	Date of mailing of the international search report 19/08/2003
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel (+31-70) 340-2040, Tx 31 651 epo nl, Fax (+31-70) 340-3016	Authorized officer VON EGGELKRAUT, S

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/03916

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0025746	A	11-05-2000	IT MI982364 A1 03-05-2000
		AU 762367 B2	26-06-2003
		AU 1266600 A	22-05-2000
		BR 9915251 A	30-10-2001
		CA 2349268 A1	11-05-2000
		CN 1335766 T	13-02-2002
		CZ 20011541 A3	12-09-2001
		WO 0025746 A2	11-05-2000
		EP 1126823 A2	29-08-2001
		HU 0104230 A2	28-03-2002
		JP 2002528484 T	03-09-2002
		NO 20012175 A	02-07-2001
		NZ 511305 A	25-07-2003
		PL 347908 A1	22-04-2002
		SK 5932001 A3	06-11-2001
		TR 200101209 T2	21-09-2001
		US 6464958 B1	15-10-2002